

Computing Two Types of Geometric-Arithmetic Indices of Some Benzenoid Graphs

AMIR LOGHMAN* AND MAHBOUBEH SAHELI

Department of Mathematics, Payame Noor Universtiy,

PO BOX 19395-3697 Tehran, I. R. IRAN.

ABSTRACT. The geometric-arithmetic index is a topological index was defined as $GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v}$, where d_u denotes the degree of vertex u in G . By replacing $\delta_u = \sum_{v \sim u} d_v$ instead of d_u in $GA(G)$, we have a new version of this index that defined as $GA_5(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\delta_u \delta_v}}{\delta_u + \delta_v}$. In this paper, we present exact formulas of these indices for some benzenoid graphs.

Keywords: benzenoid graph, geometric-arithmetic index, GA_5 index.

1. INTRODUCTION

Let G be a simple connected graph. A graph is a collection of points and lines connecting a subset of them. The points and lines of a graph also called vertices and edges of the graph, respectively. In chemical graph theory, the vertices and edges of a graph also correspond to the atoms and bonds of the molecular graph, respectively. If e is an edge/bond of G , connecting the vertices/atoms u and v , then we write

*Corresponding author. (E-mail: Amirloghman@pnu.ac.ir)

$e = uv$ say " u and v are adjacent". Chemical graph theory is an important branch of graph theory and Mathematical chemistry, which applies graph theory to mathematical modeling of chemical phenomena [7, 9, 10, 12]. A chemical topological index is a numeric quantity from the structural graph of a molecule and is invariant on the automorphism of the graph. The concept of geometric-arithmetic indices was introduced in the chemical graph theory. These indices generally are defined as:

$$GA_{general}(G) = \sum_{uv \in E(G)} \frac{2\sqrt{Q_u Q_v}}{Q_u + Q_v},$$

where Q_u is some quantity that in a unique manner can be associated with the vertex u of graph G . The first member of this class was considered by Vukičević and Furtula [11], by setting Q_u to be d_u , we have:

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v},$$

in which d_u denotes the degree of vertex u in G , i.e., the number of its neighbors in G . The second member of this class was considered by Fath-Tabar et al. [2], by setting Q_u to be the number n_u of vertices of G lying closer to the vertex u than to the vertex v for the edge uv of the graph G ,

$$GA_2(G) = \sum_{uv \in E(G)} \frac{2\sqrt{n_u n_v}}{n_u + n_v}.$$

The third member of this class was considered by Bo Zhou et al. [14], by setting Q_u to be the number m_u of edges of G lying closer to the vertex u than to the vertex v for the edge uv of the graph G ,

$$GA_3(G) = \sum_{uv \in E(G)} \frac{2\sqrt{m_u m_v}}{m_u + m_v}.$$

The fourth and fifth members of this class was considered by Ghorbani et al. [3,4], by setting Q_u to be ε_u , the eccentricity of vertex u (the largest distance between u and any other vertex of graphs) and δ_u , the summation of degree of neighbors of vertex u ,

$$GA_4(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\varepsilon_u \varepsilon_v}}{\varepsilon_u + \varepsilon_v} \text{ and } GA_5(G) = \sum_{uv \in E(G)} \frac{2\sqrt{\delta_u \delta_v}}{\delta_u + \delta_v}.$$

For a comprehensive survey of the mathematical properties and chemical properties of these indices see papers series and books [1, 5, 6, 8, 13].

A benzenoid system is a connected graph obtained by arranging congruent regular hexagons in a plane, so that two hexagons are either disjoint or have a common edge. Benzenoid graphs are simple, plane, and bipartite. The vertices lying on the border of the unbounded face of a benzenoid graphs are called external and other vertices, if present, are called internal. In this paper, we focus on the first and fifth geometric-arithmetic indices are compute them for some benzenoid graphs.

2. MAIN RESULTS

In this section, we compute first and fifth geometric-arithmetic indices for some benzenoid graphs as shown in Figure 1.




Figure 1: The benzenoid graphs T_4 , R_4 and X_4 from left to right.

The first class of benzenoid graphs we consider is triangular benzenoids such as shown in Figure 1. We denote this graph by T_n in which n is the number of hexagons in the base of graph. Obviously, the total number of hexagons in T_n is $n(n+1)/2$. Also T_n has $|V(T_n)| = n^2 + 4n + 1$ vertices and $|E(T_n)| = \frac{3}{2}n(n+3)$ edges.

Theorem 1. For graph, T_n we have:

$$GA(T_n) = \frac{3}{2}n(n-1) + 6 + \frac{12\sqrt{6}}{5}(n-1),$$

$$GA_5(T_n) = \frac{3}{2}n(n-3) + 3 + \sqrt{35} + \frac{8\sqrt{5}}{3} + \frac{12\sqrt{42}}{13}(n-2) + \frac{9\sqrt{7}}{8}(n-1).$$

Proof. Let m_{ij} is an edge that connects a vertex of degree i to a vertex of degree j . So in graph T_n for external vertices we have $|m_{22}|=6$, $|m_{23}|=6(n-1)$ and for internal vertices that all of them are from degree 3, $|m_{33}|=\frac{3}{2}n(n-1)$, then we have:

$$GA(T_n) = \frac{3}{2}n(n-1) + 6 + \frac{12\sqrt{6}}{5}(n-1).$$

For compute $GA_5(T_n)$, we partition the edges of graph in 5 subsets. All of m_{22} edges have a vertex u with $\delta(u)=5$ and a vertex v with $\delta(v)=6$. In m_{23} edges, we consider two cases. One set is contain edges with $\delta(u)=5$ and $\delta(v)=7$, that these edges have a common vertex with a m_{22} . So the number of these edges are equal to 6. Other edges with two external vertices have $\delta(u)=6$ and $\delta(v)=7$, that the number of these edges are equal to $6(n-2)$. Edges with an external vertex and an internal vertex are equal to $3(n-1)$ and have $\delta(u)=7$ and $\delta(v)=9$. Finally the number of edges with two internal vertices are $\frac{3}{2}n(n-3)+3$ and $\delta(u)=\delta(v)=9$. Then we have:

$$GA_5(T_n) = \frac{3}{2}n(n-3) + 3 + \sqrt{35} + \frac{8\sqrt{5}}{3} + \frac{12\sqrt{42}}{13}(n-2) + \frac{9\sqrt{7}}{8}(n-1).$$

Which completes the proof.

A benzenoid rhombus R_n is obtained from two copies of a triangular benzenoid T_n by identifying hexagons in one of their base rows that is shown in Figure 1. Consequently, $|V(R_n)|=2n(n+2)$ and $|E(R_n)|=3n^2+4n-1$.

Theorem 2. For graph R_n we have:

$$GA(R_n) = 3n^2 - 4n + 7 + \frac{16\sqrt{6}}{5}(n-1),$$

$$GA_5(R_n) = 3n^2 - 8n + 7 + \frac{8\sqrt{20}}{9} + \frac{4\sqrt{35}}{3} + \frac{16\sqrt{42}}{13}(n-2) + \frac{\sqrt{63}}{2}(n-1).$$

Proof. The proof is similar to theorem 1.

Third benzenoid graph that we consider is benzenoid hourglass. A benzenoid hourglass X_n is obtained from two copies of T_n by overlapping their extremal hexagons in the way shown in Figure 1. The number of vertices and edges of X_n is given by $|V(X_n)|=2(n^2+4n-2)$ and $|E(X_n)|=3n^2+9n-4$.

Theorem 3. For graph X_n we have:

$$GA(X_n) = 3n^2 - 3n + 12 + \frac{8\sqrt{6}}{5}(3n - 4),$$

$$GA_5(X_n) = 3n^2 + 9n + 10 + \frac{16\sqrt{20}}{9} + \frac{16\sqrt{35}}{12} + \frac{24\sqrt{42}}{13}(n - 2) + \frac{12\sqrt{63}}{16}(n - 1).$$

Proof. The proof is similar to Theorem 1.

In Table I, we compute $GA(G)$ and $GA_5(G)$ for three graphs in Figure 1, ($2 \leq n \leq 7$).

n	$GA(T_n)$	$GA(R_n)$	$GA(X_n)$	n	$GA_5(T_n)$	$GA_5(R_n)$	$GA_5(X_n)$
2	14.879	18.838	25.838	2	14.856	18.832	61.792
3	26.758	37.677	49.596	3	26.814	37.776	103.71
4	41.636	62.515	79.354	4	41.774	62.721	151.63
5	59.515	93.354	115.11	5	59.731	93.667	205.54
6	80.394	130.19	156.87	6	80.691	130.61	265.46
7	104.27	173.03	204.63	7	331.38	173.56	331.38

Table I.

Corollary 4. For benzenoid graphs in Figure 1 we have the following statements:

$$GA(R_n) - GA(T_n) \approx 1.5n^2 - 0.5404n - 0.9596,$$

$$GA(X_n) - GA(T_n) \approx 1.5n^2 + 4.3788n - 3.798,$$

$$GA(X_n) - GA(R_n) \approx 4.9192n - 2.8384,$$

$$GA(X_n) - 2GA(T_n) \approx -3.9192,$$

$$GA_5(T_n) - GA(T_n) \approx 0.0801n - 0.1834,$$

$$GA_5(R_n) - GA(R_n) \approx 0.1074n - 0.2193,$$

$$GA_5(X_n) - GA(X_n) \approx 18.160n - 0.366.$$

Corollary 5. Let G_n be one of the benzenoid graphs T_n, R_n or X_n , we have:

$$\lim_{n \rightarrow \infty} \frac{GA(G_n)}{GA(G_{n-1})} = 1.$$

REFERENCES

- [1] A. R. Ashrafi and H. Shabani, GA index and Zagreb indices of nanocones, *Optoelectron. Adv. Mater. Rapid Comm.*, 4(11)(2010) 1874-1876.
- [2] G. H. Fath-Tabar, B. Furtula and I. Gutman, A new geometric-arithmetic index, *J. Math. Chem.*, 47 (2010) 477-486.
- [3] M. Ghorbani and A. Khaki, A note on the fourth version of geometric-arithmetic index, *Optoelectron. Adv. Mater. Rapid Comm.*, 4 (2010) 2212-2215.
- [4] A. Graovać, M. Ghorbani and M. A. Hosseinzadeh, Computing fifth geometric arithmetic index for nanostar dendrimers, *Journal of Mathematical Nanoscience*, 1(2011) 33-42.
- [5] A. Iranmanesh and M. Zeraatkar, Computing GA index for some nanotubes, *Optoelectron. Adv. Mater. Rapid Comm.*, 4(11)(2010) 1852-1855.
- [6] A. Khaksar, M. Ghorbani and H. R. Maimani, On atom bond connectivity and GA indices of nanocones, *Optoelectron. Adv. Mater. Rapid Comm.*, 4(11)(2010) 1868-1870.
- [7] M. Randić, On characterization of molecular branching. *J. Am. Chem. Soc.*, 97 (1975) 6609-6615.
- [8] H. Shabani and A. R. Ashrafi, Computing the GA index of nanotubes and nanotori, *Optoelectron. Adv. Mater. Rapid Comm.*, 4(11)(2010) 1860-1862.
- [9] R. Todeschini, V. Consonni, *Handbook of Molecular Descriptors*, Wiley-VCH, Weinheim, 2000.
- [10] N. Trinajstić, *Chemical Graph Theory*, CRC Press, Boca Raton, FL, 1992.

- [11] D.Vukičević and B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, *J. Math. Chem.*, 46 (2009) 1369-1376.
- [12] H. Wiener, Structural determination of paraffin boiling points, *J. Am. Chem. Soc.*, 69 (1947) 17-20.
- [13] L. Xiao, S. Chen, Z. Guo and Q. Chen, The geometric-arithmetic index of Benzenoid Systems and Phenylenes, *Int. J. Contemp. Math. Sciences*. 5 (45) (2010) 2225-2230.
- [14] B. Zhou, I. Gutman, B. Furtula and Z. Du, On two types of geometric arithmetic index, *Chem. Phys. Lett.*, 482 (2009) 153-155.